

University of Science and Technology, Beijung
Optical Material and Device Lab

2022 SPRING FESTIVAL
BEIJING
CRYSTALLOGRAPHY
SCHOOL



February 1 - 14, Beijing 2022



SYMMETRY RELATIONS OF SPACE GROUPS

Mois I. Aroyo
Universidad del País Vasco, Bilbao, Spain



Universidad
del País Vasco

Euskal Herriko
Unibertsitatea

MAXIMAL SUBGROUPS OF SPACE GROUPS

Subgroups: Some basic results (summary)

Subgroup $H < G$

1. $H = \{e, h_1, h_2, \dots, h_k\} \subset G$
2. H satisfies the group axioms of G

Proper subgroups $H < G$, and
trivial subgroup: $\{e\}$, G

Index of the subgroup H in G : $[i] = |G|/|H|$
 $(\text{order of } G)/(\text{order of } H)$

Maximal subgroup H of G
NO subgroup Z exists such that:
 $H < Z < G$

Coset decomposition $G:H$

Group-subgroup pair $H < G$

left coset
decomposition

$$G = H + g_2H + \dots + g_mH, \quad g_i \notin H,$$

m=index of H in G

right coset
decomposition

$$G = H + Hg_2 + \dots + Hg_m, \quad g_i \notin H$$

m=index of H in G

Normal
subgroups

$$Hg_j = g_jH, \text{ for all } g_j = I, \dots, [i]$$

Conjugate subgroups

Conjugate subgroups

Let $H_1 < G, H_2 < G$

then, $H_1 \sim H_2$, if $\exists g \in G: g^{-1}H_1g = H_2$

(i) Classes of conjugate subgroups: $L(H)$

(ii) If $H_1 \sim H_2$, then $H_1 \cong H_2$

(iii) $|L(H)|$ is a divisor of $|G|/|H|$


Normal subgroup

$H \triangleleft G$, if $g^{-1}Hg = H$, for $\forall g \in G$


Example

Subgroups of point groups

Molecule of pentacene




Subgroups of mm2



Subgroup graph

Index



MAXIMAL SUBGROUPS OF SPACE GROUPS

I. MAXIMAL
TRANSLATIONENGLEICHE
SUBGROUPS

Subgroups of Space groups

Coset decomposition $G:T_G$

$(I, 0)$	(W_2, w_2)	...	(W_m, w_m)	...	(W_i, w_i)
(I, t_1)	$(W_2, w_2 + t_1)$...	$(W_m, w_m + t_1)$...	$(W_i, w_i + t_1)$
(I, t_2)	$(W_2, w_2 + t_2)$...	$(W_m, w_m + t_2)$...	$(W_i, w_i + t_2)$
...
(I, t_j)	$(W_2, w_2 + t_j)$...	$(W_m, w_m + t_j)$...	$(W_i, w_i + t_j)$
...

Factor group G/T_G


isomorphic to the point group P_G of G

Point group $P_G = \{I, W_2, W_3, \dots, W_i\}$

Example: P12/m1

Coset decomposition $G:T_G$


Factor group $G/T_G \approx P_G$



inversion centres (\bar{I}, t) :

$$P_G = \{ I, 2, \bar{I}, m \}$$

T_G	$T_G 2$	$T_G \bar{I}$	$T_G m$
$(I, 0)$	$(2, 0)$	$(\bar{I}, 0)$	$(m, 0)$
(I, t_I)	$(2, t_I)$	(\bar{I}, t_I)	(m, t_I)
(I, t_2)	$(2, t_2)$	(\bar{I}, t_2)	(m, t_2)
...
(I, t_j)	$(2, t_j)$	(\bar{I}, t_j)	(m, t_j)



Translationengleiche subgroups $H < G$:
***t*-subgroups**

$$\left\{ \begin{array}{l} T_H = T_G \\ P_H < P_G \end{array} \right.$$

Example: $P12/m1$

Coset decomposition
 $G : T_G$

***t*-subgroups:**

$$H_1 = T_G \cup T_{G2}$$

$P121$

T_G	T_{G2}
$(1,0)$	$(2,0)$
$(1,t_1)$	$(2,t_1)$
$(1,t_2)$	$(2,t_2)$
...	...
$(1,t_j)$	$(2,t_j)$
...	...


$$P\bar{1} = H_2 = T_G \cup T_{G\bar{1}}$$

$T_G\bar{1}$	T_Gm
$(\bar{1},0)$	$(m,0)$
$(\bar{1},t_1)$	(m,t_1)
$(\bar{1},t_2)$	(m,t_2)
...	...
$(\bar{1},t_j)$	(m,t_j)
...	...

$$H_3 = T_G \cup T_Gm$$

Pm



Example: P12/m1




Translationengleiche
subgroups $H < G$:

$$P\bar{1} = T_G \cup T_{G\bar{1}}$$

$$P12\bar{1} = T_G \cup T_{G2}$$



Example: P12/m I



Subgroup diagram of point group $2/m$


Translationengleiche subgroups $H < G$:

index


[1]

[2]

[4]



Translationengleiche subgroups of space group $P2/m$

Transformation matrix: (P,p) Subgroup specification: HM symbol, [i], (P,p)

MAXIMAL SUBGROUPS OF SPACE GROUPS

II. MAXIMAL
KLASSENGLAICHE
SUBGROUPS

Klassengleiche subgroups $H < G$:

Example: P I

$$t = ua + vb + wc$$

Coset decomposition

$$T_e = \{t(u=2n, v, w)\}$$

$$t_a(a, 0, 0)$$

isomorphic k -subgroups:

$$P I(2a, b, c)$$

Subgroups of space groups

$$\left\{ \begin{array}{l} T_H < T_G \\ P_H = P_G \end{array} \right.$$

T_e	$T_e t_a$
$(l, 0)$	(l, t_a)
(l, t_1)	$(l, t_1 + t_a)$
(l, t_2)	$(l, t_2 + t_a)$
...	...
(l, t_j)	$(l, t_j + t_a)$
...	...

$$H = T_e$$

Klassengleiche subgroups $H < G$:

Example: PI

$$t = u\mathbf{a} + v\mathbf{b} + w\mathbf{c}$$

Coset decomposition

$$PI = T_e + T_e t_a$$

$$T_e = \{t(u=2n, v, w)\}$$

Isomorphic k -subgroup:

$$PI(2\mathbf{a}, \mathbf{b}, \mathbf{c})$$

Series of isomorphic k -subgroups:

$$PI(p\mathbf{a}, \mathbf{b}, \mathbf{c}): p > l, \text{ prime}$$

$$PI(a, q\mathbf{b}, \mathbf{c}): q > l, \text{ prime}$$

... etc.

Subgroups of space groups

$$\begin{cases} T_H < T_G \\ P_H = P_G \end{cases}$$

$$H = T_e \quad t_a(l, 0, 0)$$

T_e	$T_e t_a$
$(l, 0)$	(l, t_a)
(l, t_1)	$(l, t_1 + t_a)$
(l, t_2)	$(l, t_2 + t_a)$
...	...
(l, t_j)	$(l, t_j + t_a)$
...	...

INFINITE number of maximal isomorphic subgroups

Example: P- I

Series of maximal isomorphic subgroups

 $P\bar{1}$

No. 2

 $P\bar{1}$

• Series of maximal isomorphic subgroups

$[p] \mathbf{a}' = p\mathbf{a}, \mathbf{b}' = q\mathbf{a} + \mathbf{b}, \mathbf{c}' = r\mathbf{a} + \mathbf{c}$

 $P\bar{1}$ (2)

$\langle 2 + (2u, 0, 0) \rangle$

$p > 2; 0 \leq q < p; 0 \leq r < p; 0 \leq u < p$

p conjugate subgroups for each triplet of q, r , and prime p

$p\mathbf{a}, q\mathbf{a} + \mathbf{b}, r\mathbf{a} + \mathbf{c}$

$u, 0, 0$

$[p] \mathbf{b}' = p\mathbf{b}, \mathbf{c}' = q\mathbf{b} + \mathbf{c}$

 $P\bar{1}$ (2)

$\langle 2 + (0, 2u, 0) \rangle$

$p > 2; 0 \leq q < p; 0 \leq u < p$

p conjugate subgroups for each pair of q and prime p

$\mathbf{a}, p\mathbf{b}, q\mathbf{b} + \mathbf{c}$

$0, u, 0$

$[p] \mathbf{c}' = p\mathbf{c}$

 $P\bar{1}$ (2)

$\langle 2 + (0, 0, 2u) \rangle$

$p > 2; 0 \leq u < p$

p conjugate subgroups for the prime p

$\mathbf{a}, \mathbf{b}, p\mathbf{c}$

$0, 0, u$

Klassengleiche subgroups $H < G$: **k -subgroups**

$$\left\{ \begin{array}{l} T_H < T_G \\ P_H = P_G \end{array} \right.$$

Example: $P\bar{1}2/m\bar{1}$

Klassengleiche
subgroup
 $P2/m > P2/m(2\mathbf{b})$
isomorphic

$$P2/m(2\mathbf{b}) = T_{2\mathbf{b}} P_G$$

non-isomorphic
 k -subgroups:

Coset decomposition $G:T_G$

T_G $T_G 2$ $T_G \bar{1}$ $T_G m$

$(1, 0)$	$(2, 0)$	$(\bar{1}, 0)$	$(m, 0)$
$(1, t_1)$	$(2, t_1)$	$(\bar{1}, t_1)$	(m, t_1)
$(1, t_2)$	$(2, t_2)$	$(\bar{1}, t_2)$	(m, t_2)
...
$(1, t_j)$	$(2, t_j)$	$(\bar{1}, t_j)$	(m, t_j)
...

$P2/m > P2\bar{1}/m(2\mathbf{b})$

$P2\bar{1}/m(2\mathbf{b}) = T_{2\mathbf{b}} \cup T_{2\mathbf{b}t_{\mathbf{b}}2} \cup T_{2\mathbf{b}\bar{1}} \cup T_{2\mathbf{b}t_{\mathbf{b}}m}$

$$t_{\mathbf{b}} = (0, 1, 0)$$

Klassengleiche subgroups $H < G$:
non-isomorphic

$$\left\{ \begin{array}{l} T_H < T_G \\ P_H = P_G \end{array} \right.$$

Example: C2

Coset decomposition


$$C2 = T_c + T_{c2}$$

$$(T_i + T_{it_c})$$

$$\begin{aligned} t_i &= \text{integer} \\ t_c &= 1/2, 1/2, 0 \end{aligned}$$

non-isomorphic
k-subgroups:

$$\begin{aligned} H_1 &= T_i \cup T_{i2} \\ &P2 \end{aligned}$$



Example: P4mm

Maximal subgroups of space groups

C_{4v}^1

$P4mm$

No. 99

$P4mm$

I Maximal *translationengleiche* subgroups

[2] $P411$ (75, $P4$)	1; 2; 3; 4
[2] $P21m$ (35, $Cmm2$)	1; 2; 7; 8
[2] $P2m1$ (25, $Pmm2$)	1; 2; 5; 6

$\mathbf{a} - \mathbf{b}, \mathbf{a} + \mathbf{b}, \mathbf{c}$



II Maximal *klassengleiche* subgroups

- Enlarged unit cell

[2] $\mathbf{c}' = 2\mathbf{c}$		
$P4_2mc$ (105)	$\langle 2; 5; 3 + (0, 0, 1) \rangle$	$\mathbf{a}, \mathbf{b}, 2\mathbf{c}$
$P4cc$ (103)	$\langle 2; 3; 5 + (0, 0, 1) \rangle$	$\mathbf{a}, \mathbf{b}, 2\mathbf{c}$
$P4_2cm$ (101)	$\langle 2; (3; 5) + (0, 0, 1) \rangle$	$\mathbf{a}, \mathbf{b}, 2\mathbf{c}$
$P4mm$ (99)	$\langle 2; 3; 5 \rangle$	$\mathbf{a}, \mathbf{b}, 2\mathbf{c}$

- Series of maximal isomorphic subgroups


[p] $\mathbf{c}' = p\mathbf{c}$			
$P4mm$ (99)	$\langle 2; 3; 5 \rangle$ $p > 1$ no conjugate subgroups	$\mathbf{a}, \mathbf{b}, p\mathbf{c}$	
$[p^2] \mathbf{a}' = p\mathbf{a}, \mathbf{b}' = p\mathbf{b}$	$\langle 2 + (2u, 2v, 0); 3 + (u + v, -u + v, 0); 5 + (0, 2v, 0) \rangle$ $p > 2; 0 \leq u < p; 0 \leq v < p$ p^2 conjugate subgroups for the prime p	$p\mathbf{a}, p\mathbf{b}, \mathbf{c}$	$u, v, 0$



GENERAL SUBGROUPS OF SPACE GROUPS

General subgroups $H < G$:

Graph of maximal subgroups



Group-subgroup pair

$$G > \mathcal{H} : G, \mathcal{H}, [i], (P, p)$$

Pairs: group - maximal subgroup

$$\mathcal{Z}_k > \mathcal{Z}_{k+1}, (P, p)_k$$

$$(P, p) = \prod_{k=1}^n (P, p)_k$$

General subgroups $H < G$:

$$\left\{ \begin{array}{l} T_H < T_G \\ P_H < P_G \end{array} \right.$$

Theorem Hermann, 1929:

For each pair $G > H$, there exists a uniquely defined intermediate subgroup M , $G \geq M \geq H$, such that:

M is a *t*-subgroup of G

H is a *k*-subgroup of M

$$[i] = [i_P] \cdot [i_L]$$




Corollary

A maximal subgroup is either a *t*- or *k*-subgroup

Subgroups of Space groups

Coset decomposition $G:T_G$



	(l,0)	(W ₂ ,w ₂)	...	(W _m ,w _m)	...	(W _i ,w _i)
	(l,t ₁)	(W ₂ ,w ₂ +t ₁)	...	(W _m ,w _m +t ₁)	...	(W _i ,w _i +t ₁)
	(l,t ₂)	(W ₂ ,w ₂ +t ₂)	...	(W _m ,w _m +t ₂)	...	(W _i ,w _i +t ₂)
...
	(l,t _j)	(W ₂ ,w ₂ +t _j)	...	(W _m ,w _m +t _j)	...	(W _i ,w _i +t _j)
...

Factor group G/T_G

isomorphic to the point group P_G of G



Point group $P_G = \{l, W_2, W_3, \dots, W_i\}$

Example:

P4mm (a,b,c) > Pmm2 (2a,b,c)

[i]=4

$$[i] = [i_P] \cdot [i_L]$$



DOMAIN-STRUCTURE ANALYSIS (INITIAL STEPS)

PROBLEM:

Domain-structure analysis

$$G \xrightarrow{[i]} H$$

number of domain states

twins and antiphase domains

twinning operation

symmetry groups of the domain states; multiplicity and degeneracy

Phase transitions domain structures

Homogeneous
(parent) phase



Deformed
(daughter) phase
Domain structure

When a **crystal homogeneous** in the parent (prototypic, high-symmetry) phase undergoes a phase transition into a low-symmetry phase (ferroic, if the point-group symmetry is lowered) then this **daughter** phase is almost always formed as a **non-homogeneous** structure consisting of **homogeneous regions** called **domains**

Domain

A connected homogeneous part of a domain structure or of a twinned crystal is called a **domain**. Each domain is a single crystal.

Different domains can exhibit different **tensor properties**, different **diffraction patterns** and can differ in other physical properties.

optical observation of domain structure




Fig. 3.4.1.1. Domain structure of tetragonal barium titanate (BaTiO_3). A thin section of barium titanate ceramic observed at room temperature in a polarized-light microscope (transmitted light, crossed polarizers). Courtesy of U. Täffner, Max-Planck-Institut für Metallforschung, Stuttgart. Different colours correspond to different ferroelastic domain states, connected areas of the same colour are ferroelastic domains and sharp boundaries between these areas are domain walls. Areas of continuously changing colour correspond to gradually changing thickness of wedge-shaped domains. An average distance between parallel ferroelastic domain walls is of the order of 1–10 μm .

Powerful high-resolution electron microscopy (HREM)




Fig. 3.4.1.2. Domain structure of a BaGa_2O_4 crystal seen by high-resolution transmission electron microscopy. Parallel rows are atomic layers. Different directions correspond to different ferroelastic domain states of domains, connected areas with parallel layers are different ferroelastic domains and boundaries between these areas are ferroelastic domain walls. Courtesy of H. Lemmens, EMAT, University of Antwerp.

Phase transitions domain structures

Homogeneous
(parent) phase



Deformed
(daughter) phase
Domain structure

Domains

The **number** of such crystals **is not limited**; they differ in their locations in space, in their orientations, in their shapes and in their space groups but all belong to the **same space-group type of H**.

Domain states

The domains belong to a finite (small) number of **domain states**.

Two domains belong to the same *domain state* if their crystal patterns are identical, i.e. if they occupy different regions of space that are part of the **same crystal pattern**.


The **number of domain states** which are observed after a phase transition is limited and **determined by** the group-subgroup relations of the **space groups G and H**.

Symmetry Reduction

initial phase



daughter phase



symmetry of
a square

symmetry of
a rectangle

two possible
orientations

SUBGROUPS CALCULATIONS: HERMANN

Hermann, 1929:

For each pair $G > \mathcal{H}$, index $[i]$, there exists a uniquely defined intermediate subgroup \mathcal{M} , $G \geq \mathcal{M} \geq \mathcal{H}$, such that:


\mathcal{M} is a *t*-subgroup of G

\mathcal{H} is a *k*-subgroup of \mathcal{M}

with $[i] = [i_P] \cdot [i_L]$

$$i_P = P_G / P_H$$

$$i_L = Z_{H,P} / Z_{G,P} = V_{H,P} / V_{G,P}$$



twins

antiphase

EXAMPLE

Lead vanadate $\text{Pb}_3(\text{VO}_4)_2$

Index [i] for a group-subgroup pair $G>H$

$\mathcal{R}-3m$

$$i_P = P_G / P_H$$

$$[i_P] = 3$$

$C2/m$


$$i_L = Z_{H,p} / Z_{G,p}$$

$$[i_L] = 2$$

$P2_1/c$

High-symmetry phase $R-3m$

166	5.6748	5.6748	20.3784	90	90	120	Z_{G,p}=1	 P_G =12
5								
Pb	1	3a	0.000000		0.000000		0.000000	
Pb	2	6c	0.000000		0.000000		0.207100	
PV	3	6c	0.000000		0.000000		0.388400	
0	4	6c	0.000000		0.000000		0.324000	
0	5	18i	0.842400		0.157600		0.430100	



Low-symmetry phase $P2_1/c$

14	7.5075	6.0493	9.4814	90.	115.162	90.
7						
Pb	1	2a	0 0 0			
Pb	2	4e	0.3835	0.5815	0.2879	
PV	1	4e	0.2071	0.0143	0.3999	
0	1	4e	0.2872	0.2559	0.0159	
0	2	4e	0.2598	0.7979	0.0216	
0	3	4e	0.3194	0.9784	0.2823	
0	4	4e	0.0335	0.5431	0.2091	

$$|P_H|=?$$


$$Z_{H,p}=?$$

Pb₃(VO₄)₂: Ferroelastic Domains in P2₁/c phase

Group-Subgroup Lattice

Maximal-subgroup graph

SUBGROUPGRAPH



number of domain states = index [i] = [i_P].[i_L] = 6


number of ferroelastic domain states: i_P = 12:4 = 3

number of different subgroups P2₁/c: 3

Problem:

CLASSIFICATION OF DOMAINS

HERMANN



EXERCISES

(Problem 2.6.3)

- (A) High symmetry phase: P2/m
Low symmetry phase: PI, small unit-cell deformation
How many and what kind of domain states?

Hint: Determine the index $[i]=[i_P].[i_L]$

- (B) High symmetry phase: P2/m
Low symmetry phase: PI, duplication of the unit cell

How many and what kind of domain states?

- (C) High symmetry phase: P4mm
Low symmetry phase: P2, index 8

How many and what kind of domain states?

- (D) High symmetry phase: P4₂bc
Low symmetry phase: P2₁, index 8

How many and what kind of domain states?

EXERCISES

(Problem 2.6.4)


At high temperatures, BiTiO_3 has the cubic perovskite structure, space group Pm-3m (No. 221). Upon cooling, it distorts to three slightly deformed structures, all three being ferroelectric, with space groups $\text{P}4\text{mm}$ (No. 99), $\text{Amm}2$ (No. 38) and $\text{R}3\text{m}$ (No. 160). Can we expect twinned crystals of the low symmetry forms? If so, how many and what kind of domain states could occur?

Hint: The program INDEX could be useful


Problem 2.6.4

SOLUTION


(a) cubic
($T > 120^\circ\text{C}$)




(b) tetragonal
($120^\circ\text{C} > T > 5^\circ\text{C}$)



(c) orthorhombic
($5^\circ\text{C} > T > -90^\circ\text{C}$)




(d) rhombohedral
($T < -90^\circ\text{C}$)



Unit cells of the four phases of BaTiO_3

BaTiO₃: Ferroelectric Domains in P4mm phase



Maximal-subgroup graph



$$\text{index } [i] = i_p = 48 : 8 = 6$$

number of ferroelectric domain states: 6

number of different subgroups P4mm: 3



Domain-structure analysis: Twinning operation

Coset decomposition of G:H

left: $G>H, G=H+(V_2, v_2)H + \dots + (V_n, v_n)H$

right: $G>H, G=H+H(W_2, w_2) + \dots + H(W_n, w_n)$

Please, enter the sequential numbers of group and subgroup as given in International Tables for Crystallography, Vol. A:

Enter supergroup number (G) or [choose it](#):

221

Enter subgroup number (H) or [choose it](#):

99

Please, define the [transformation](#) that relates the group and the subgroup bases.

Enter transformation matrix :

Rotational part			Origin Shift
1	0	0	0
0	1	0	0
0	0	1	0

Decomposition:

left right

BaTiO₃: Ferroelectric Domains in P4mm phase

Twinning operations

Coset decomposition: Pm $\bar{3}$ m : P4_zmm, index 6

Coset 1:	Coset 2:	Coset 3:	Coset 4:	Coset 5:	Coset 6:
(x, y, z)	(-x, y, -z)	(z, x, y)	(-z, -x, y)	(y, z, x)	(y, -z, -x)
(-x, -y, z)	(x, -y, -z)	(z, -x, -y)	(-z, x, -y)	(-y, z, -x)	(-y, -z, x)
(-y, x, z)	(y, x, -z)	(z, -y, x)	(-z, y, x)	(x, z, -y)	(x, -z, y)
(y, -x, z)	(-y, -x, -z)	(z, y, -x)	(-z, -y, -x)	(-x, z, y)	(-x, -z, -y)
(x, -y, z)	-x, -y, -z)	(z, x, -y)	(-z, -x, -y)	(-y, z, x)	-y, -z, -x)
(-x, y, z)	(x, y, -z)	(z, -x, y)	(-z, x, y)	(y, z, -x)	(y, -z, x)
(-y, -x, z)	(y, -x, -z)	(z, -y, -x)	(-z, y, -x)	(-x, z, -y)	(-x, -z, y)
(y, x, z)	(-y, x, -z)	(z, y, x)	(-z, -y, x)	(x, z, y)	(x, -z, -y)

coset representatives: q_i

(1,0) ($\bar{1}$,0) (3,0) ($\bar{3}$,0) (3⁻¹,0) ($\bar{3}^{-1}$,0)

polarization: $P_i = q_i P$

0
0
v

0
0
-v

v
0
0

-v
0
0


0
v
0

0
-v
0

RELATIONS BETWEEN WYCKOFF POSITIONS

Relations between Wyckoff positions


$$\mathcal{G} = \text{Pmm2} > \mathcal{H} = \text{Pm}, [i] = 2$$



$S_0, \mathcal{G} = \text{Pmm2}$

$2h$ m.. $(0,y,z)$

$2f$.m. $(x,0,z)$



$S_1, \mathcal{H} = \text{Pm}$

$2c$ | (x,y,z)

|
 b m $(x_2,0,z_2)$
|
 b m $(x_1,0,z_1)$

SYMMETRY REDUCTION

EXAMPLE

Consider the group
-subgroup pair $P4mm > Pmm2$
 $[i]=2, a'=a, b'=b, c'=c$

Determine the splitting schemes for WPs 1a, 1b, 2c, 4d, 4e

group $P4mm$

Generators selected (1); $t(1,0,0)$; $t(0,1,0)$; $t(0,0,1)$; (2); (3); (5)

Positions

Multiplicity,
Wyckoff letter,
Site symmetry

		Coordinates			
8	<i>g</i>	1	(1) x, y, z	(2) \bar{x}, \bar{y}, z	(3) \bar{y}, x, z
			(5) x, \bar{y}, z	(6) \bar{x}, y, z	(7) \bar{y}, \bar{x}, z
					(8) y, x, z
4	<i>f</i>	. <i>m</i> .	$x, \frac{1}{2}, z$	$\bar{x}, \frac{1}{2}, z$	$\frac{1}{2}, x, z$
4	<i>e</i>	. <i>m</i> .	$x, 0, z$	$\bar{x}, 0, z$	$0, x, z$
4	<i>d</i>	. . <i>m</i>	x, x, z	\bar{x}, \bar{x}, z	\bar{x}, x, z
2	<i>c</i>	2 <i>m m</i> .	$\frac{1}{2}, 0, z$	$0, \frac{1}{2}, z$	
1	<i>b</i>	4 <i>m m</i>	$\frac{1}{2}, \frac{1}{2}, z$		
1	<i>a</i>	4 <i>m m</i>	$0, 0, z$		

subgroup $Pmm2$

Generators selected (1); $t(1,0,0)$; $t(0,1,0)$; $t(0,0,1)$; (2); (3)

Positions

Multiplicity,
Wyckoff letter,
Site symmetry


4	<i>i</i>	1	(1) x, y, z	(2) \bar{x}, \bar{y}, z	(3) x, \bar{y}, z	(4) \bar{x}, y, z
2	<i>h</i>	<i>m</i> ..		$\frac{1}{2}, y, z$	$\frac{1}{2}, \bar{y}, z$	
2	<i>g</i>	<i>m</i> ..		$0, y, z$	$0, \bar{y}, z$	
2	<i>f</i>	. <i>m</i> .		$x, \frac{1}{2}, z$	$\bar{x}, \frac{1}{2}, z$	
2	<i>e</i>	. <i>m</i> .		$x, 0, z$	$\bar{x}, 0, z$	
1	<i>d</i>	<i>m m</i> 2		$\frac{1}{2}, \frac{1}{2}, z$		
1	<i>c</i>	<i>m m</i> 2		$\frac{1}{2}, 0, z$		
1	<i>b</i>	<i>m m</i> 2		$0, \frac{1}{2}, z$		
1	<i>a</i>	<i>m m</i> 2		$0, 0, z$		

EXAMPLE


Group-subgroup pair
 $P4mm > Pmm2$, $[i]=2$

$$a'=a, b'=b, c'=c$$

$P4mm$



$Pmm2$



2c 2mm. I/2 0 z
0 I/2 z ★



★ I/2 0 z Ic mm2
★ 0 I/2 z' Ib mm2

Data on Relations between Wyckoff Positions in *International Tables for Crystallography*, Vol.AI

C_{4v}^1

No. 99

$P4mm$

Axes	Coordinates	Wyckoff positions						
		1a	1b	2c	4d	4e	4f	8g
I Maximal <i>translationengleiche</i> subgroups								
[2] $P4$ (75)		1a	1b	2c	4d	4d	4d	$2 \times 4d$
[2] $Pmm2$ (25)		1a	1d	1b; 1c	4i	2e; 2g	2f; 2h	$2 \times 4i$
[2] $Cmm2$ (35)	$\mathbf{a}-\mathbf{b}$, $\frac{1}{2}(x-y), \frac{1}{2}(x+y), z$ $\mathbf{a}+\mathbf{b}, \mathbf{c}$	2a	2b	4c	4d; 4e	8f	8f	$2 \times 8f$

II Maximal *klassengleiche* subgroups Enlarged unit cell, non-isomorphic

[2] $I4cm$ (108)	$\mathbf{a}-\mathbf{b}$, $\frac{1}{2}(x-y), \frac{1}{2}(x+y), \frac{1}{2}z;$ $\mathbf{a}+\mathbf{b}, 2\mathbf{c}$ $+(0, 0, \frac{1}{2})$	4a	4b	8c	16d	16d	$2 \times 8c$	$2 \times 16d$
[2] $I4cm$ (108)	$\mathbf{a}-\mathbf{b}$, $\frac{1}{2}(x-y)+\frac{1}{2}, \frac{1}{2}(x+y), \frac{1}{2}z;$ $\mathbf{a}+\mathbf{b}, 2\mathbf{c}$ $+(0, 0, \frac{1}{2})$	4b	4a	8c	16d	$2 \times 8c$	16d	$2 \times 16d$
[2] $I4mm$ (107)	$\mathbf{a}-\mathbf{b}$, $\frac{1}{2}(x-y), \frac{1}{2}(x+y), \frac{1}{2}z;$ $\mathbf{a}+\mathbf{b}, 2\mathbf{c}$ $+(0, 0, \frac{1}{2})$	$2 \times 2a$	4b	8c	$2 \times 8d$	$2 \times 8c$	16e	$2 \times 16e$
[2] $I4mm$ (107)	$\mathbf{a}-\mathbf{b}$, $\frac{1}{2}(x-y)+\frac{1}{2}, \frac{1}{2}(x+y), \frac{1}{2}z;$ $\mathbf{a}+\mathbf{b}, 2\mathbf{c}$ $+(0, 0, \frac{1}{2})$	4b	$2 \times 2a$	8c	$2 \times 8d$	16e	$2 \times 8c$	$2 \times 16e$
[2] $P4_2mc$ (105)	$\mathbf{a}, \mathbf{b}, 2\mathbf{c}$ $x, y, \frac{1}{2}z; +(0, 0, \frac{1}{2})$	2a	2b	$2 \times 2c$	8f	$2 \times 4d$	$2 \times 4e$	$2 \times 8f$
[2] $P4cc$ (103)	$\mathbf{a}, \mathbf{b}, 2\mathbf{c}$ $x, y, \frac{1}{2}z; +(0, 0, \frac{1}{2})$	2a	2b	4c	8d	8d	8d	$2 \times 8d$
[2] $P4_2cm$ (101)	$\mathbf{a}, \mathbf{b}, 2\mathbf{c}$ $x, y, \frac{1}{2}z; +(0, 0, \frac{1}{2})$	2a	2b	4c	$2 \times 4d$	8e	8e	$2 \times 8e$
[2] $P4bm$ (100)	$\mathbf{a}-\mathbf{b}$, $\frac{1}{2}(x-y), \frac{1}{2}(x+y), z;$ $\mathbf{a}+\mathbf{b}, -\mathbf{c}$ $-(\frac{1}{2}, \frac{1}{2}, 0)$	2a	2b	4c	8d	8d	$2 \times 4c$	$2 \times 8d$

Example

Wyckoff Positions Splitting

Conventional Settings

Non conventional Settings

Please, enter the sequential numbers of group and subgroup as given in International Tables for Crystallography, Vol. A:

Enter supergroup or [choose it](#)

136

group

Enter subgroup or [choose it](#)

65

subgroup

Please, define the transformation relating the group and the subgroup bases.

(NOTE: If you don't know the transformation click [here](#) for possible workarounds)

rotational matrix:

Transformation
matrix (P,P)

1	1	0
-1	1	0
0	0	1

origin shift:

0	0	0
---	---	---

[Show group-subgroup data.](#)

Two-level input:
Choice of the
Wyckoff positions

Wyckoff Positions Splitting

136 ($P4_2/mnm$) > 65 ($Cmmm$)

Group Data Subgroup Data

16r (x, y, z)

8q (x, y, 1/2)

8p (x, y, 0)

All positions

8o (x, 0, z)

16k (x, y, z)

8n (0, y, z)

8j (x, x, z)

8m (1/4 , 1/4 , z)

8i (x, y, 0)

4l (0, 1/2 , z)

8h (0, 1/2 , z)

4k (0, 0, z)

4g (x, - x, 0)

4j (0, y, 1/2)

4f (x, x, 0)

4i (0, y, 0)

4e (0, 0, z)

4h (x, 0, 1/2)

4d (0, 1/2 , 1/4)

4g (x, 0, 0)

Wyckoff Positions Splitting

99 ($P4mm$) > 8 (Cm) [unique axis b]

Bilbao Crystallographic Server

Result from splitting

No	Wyckoff position(s)		
	Group	Subgroup	More...
1	8g	4b 4b 4b 4b	Relations
2	4f	4b 4b	Relations
3	4e	4b 4b	Relations
4	4d	4b 2a 2a	Relations
5	2c	4b	Relations
6	1b	2a	Relations
7	1a	2a	Relations

Two-level output:

Relations between coordinate triplets

Splitting of Wyckoff position 4d

Representative		Subgroup Wyckoff position		
No	group basis	subgroup basis	name[n]	
1	(x, x, z)	(0, x, z)	4b ₁	(x ₁ , y ₁ , z ₁)
2	(-x, -x, z)	(0, -x, z)		(x ₁ , -y ₁ , z ₁)
3	(x+1, x, z)	(1/2, x+1/2, z)		(x ₁ +1/2, y ₁ +1/2, z ₁)
4	(-x+1, -x, z)	(1/2, -x+1/2, z)		(x ₁ +1/2, -y ₁ +1/2, z ₁)
5	(-x, x, z)	(-x, 0, z)	2a ₁	(x ₂ , 0, z ₂)
6	(-x+1, x, z)	(-x+1/2, 1/2, z)		(x ₂ +1/2, 1/2, z ₂)
7	(x, -x, z)	(x, 0, z)	2a ₂	(x ₃ , 0, z ₃)
8	(x+1, -x, z)	(x+1/2, 1/2, z)		(x ₃ +1/2, 1/2, z ₃)


NORMALIZERS OF SPACE GROUPS

Normalizers of space groups

Normalizers $N(G)$: $g^{-1}\{G\}g = \{G\}$ { Euclidean
Affine }

Example: Pmmn


the symmetry of
symmetry



Normalizers of space groups

Normalizers $N(G)$: $g^{-1}\{G\}g = \{G\}$ { Euclidean
 Affine }

the symmetry
of symmetry




Space group: Pmmn (a,b,c)

Euclidean normalizer:
Pmmm (1/2a, 1/2b, 1/2c)

Normalizers for specialized metrics

Normalizers



Space group:
Pmmn (a,b,c), **a=b**

Euclidean normalizer for
specialized metrics:
P4/mmm (1/2a, 1/2b, 1/2c)

Applications:

- Equivalent point configurations
- Wyckoff sets
- Equivalent structure descriptions

Normalizers of space groups

E. Koch and W. Fischer

Space group \mathcal{G}			Euclidean normalizer $\mathcal{N}_{\mathcal{E}}(\mathcal{G})$	
No.	Hermann–Mauguin symbol	Cell metric	Symbol	Basis vectors
55	$Pbam$	$a \neq b$	$Pmmm$	$\frac{1}{2}\mathbf{a}, \frac{1}{2}\mathbf{b}, \frac{1}{2}\mathbf{c}$
		$a = b$	$P4/mmm$	$\frac{1}{2}\mathbf{a}, \frac{1}{2}\mathbf{b}, \frac{1}{2}\mathbf{c}$
56	$Pccn$	$a \neq b$	$Pmmm$	$\frac{1}{2}\mathbf{a}, \frac{1}{2}\mathbf{b}, \frac{1}{2}\mathbf{c}$
		$a = b$	$P4/mmm$	$\frac{1}{2}\mathbf{a}, \frac{1}{2}\mathbf{b}, \frac{1}{2}\mathbf{c}$
57	$Pbcm$		$Pmmm$	$\frac{1}{2}\mathbf{a}, \frac{1}{2}\mathbf{b}, \frac{1}{2}\mathbf{c}$
58	$Pnnm$	$a \neq b$	$Pmmm$	$\frac{1}{2}\mathbf{a}, \frac{1}{2}\mathbf{b}, \frac{1}{2}\mathbf{c}$
		$a = b$	$P4/mmm$	$\frac{1}{2}\mathbf{a}, \frac{1}{2}\mathbf{b}, \frac{1}{2}\mathbf{c}$
59	$Pmmn$ (both origins)	$a \neq b$	$Pmmm$	$\frac{1}{2}\mathbf{a}, \frac{1}{2}\mathbf{b}, \frac{1}{2}\mathbf{c}$
		$a = b$	$P4/mmm$	$\frac{1}{2}\mathbf{a}, \frac{1}{2}\mathbf{b}, \frac{1}{2}\mathbf{c}$



Example: Pmmn

Problem: Normalizers
of space groups

NORMALIZER

Normalizers of Space Groups

How to select the group

The space groups are specified by their number as given in the *International Tables for Crystallography*, Vol. A. You can give this number, if you know it, or you can choose it from the table with the space group numbers and symbols if you click on the link [\[choose it\]](#).

Please, enter the sequential number of group as given in the *International Tables for Crystallography*, Vol. A

[choose](#)

Choose:

Euclidean (general metric):
 Enhanced Euclidean (specialized metric):
 Affine:



Enhanced Euclidean normalizer (specialized metrics)


Space group:

Lattice parameters:

[Show](#)

Example NORMALIZER: Space group *Pnnm* (59)

Euclidean normalizer (general metric) of *Pmmn* (No. 59)



Space group: *Pmmn* (59)

Lattice type: oP

Cell parameters: 4 4 5 90 90 90

Angular tolerance: 0.15 degrees

Euclidean normalizer of *Pmmn* (a,b,c): *Pmmm* (1/2a,1/2b,1/2c).

Index of *Pmmn* in *Pmmm* (1/2a,1/2b,1/2c): 8 with $i_L=8$ and $i_P=1$.

Additional generators of *Pmmm* (1/2a,1/2b,1/2c) with respect to *Pmmn*.

$x+1/2, y, z$	$\begin{pmatrix} 1 & 0 & 0 & 1/2 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}$	$t(1/2, 0, 0)$
$x, y+1/2, z$	$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1/2 \\ 0 & 0 & 1 & 0 \end{pmatrix}$	$t(0, 1/2, 0)$
$x, y, z+1/2$	$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1/2 \end{pmatrix}$	$t(0, 0, 1/2)$

Cosets representatives

x, y, z
 $x+1/2, y, z$
 $x, y+1/2, z$
 $x+1/2, y+1/2, z$
 $x, y, z+1/2$
 $x+1/2, y, z+1/2$
 $x, y+1/2, z+1/2$
 $x+1/2, y+1/2, z+1/2$

SUPERGROUPS OF SPACE GROUPS

Supergroups of space groups

Definition:


The group G is a supergroup of H if H is a subgroup of G , $G \geq H$

If H is a proper subgroup of G , $H < G$, then G is a proper supergroup of H , $G > H$

If H is a maximal subgroup of G , $H < G$, then G is a minimal supergroup of H , $G > H$

Types of minimal supergroups:

translationengleiche (t-type)
klassengleiche (k-type)




ITAI data:

minimal non-isomorphic k- and t-supergroups types

The Supergroup Problem

Given a group-subgroup pair $G > H$ of index $[i]$

Determine: all $G_k > H$ of index $[i]$, $G_i \approx G$




all $G_k > H$ contain H as subgroup

$$G_k = H + Hg_2 + \dots + Hg_{ik}$$


Example: Supergroup problem

Group-subgroup pair
 $422 > 222$



How many are
the subgroups
 222 of 422 ?

Supergroups 422 of
the group 222





How many are
the supergroups
 422 of 222 ?

Example: Supergroup problem

Group-subgroup pair
 $422 > 222$

Supergroups 422 of
the group 222



$$4_z22 = 2_z2_x2_y + 4_z(2_z2_x2_y)$$

$$4_z22 = 2_z2_{+2-} + 4_z(2_z2_{+2-})$$

$$4_z22 = 222 + 4_z222$$


$$4_y22 = 222 + 4_y222$$

$$4_x22 = 222 + 4_x222$$

Example: Supergroup problem

Group-subgroup pair
 $P422 > P222$

$P422$
[2]
 $P222$



$$P422 = 222 + (222)(4,0)$$


Supergroups $P422$ of
the group $P222$

$P4_z22$ $P4_x22$ $P4_y22$
[2]
 $P222$

$$\begin{aligned} P4_z22 &= 222 + (222)(4_z,0) \\ P4_x22 &= 222 + (222)(4_x,0) \\ P4_y22 &= 222 + (222)(4_y,0) \end{aligned}$$

**Are there more
supergroups $P422$ of $P222$?**


Example: Supergroups P422 of P222



$$\mathcal{H} = \text{P222}$$

$$\mathcal{G} = \text{P422}$$

$$\text{P422} = \text{P222} + (4|\omega)\text{P222}$$



	4 en	ω	\mathcal{G}
4_z	$(0, 0, 0)$	$(0, 0, 0)$	$(\text{P422})_1$
4_y	$(0, 0, 0)$	$(0, 0, 0)$	$(\text{P422})_2$
4_x	$(0, 0, 0)$	$(0, 0, 0)$	$(\text{P422})_3$
4_z	$(\frac{1}{2}, 0, 0)$	$(\frac{1}{2}, \frac{1}{2}, 0)$	$(\text{P422})'_1$
4_y	$(\frac{1}{2}, 0, 0)$	$(\frac{1}{2}, 0, \frac{1}{2})$	$(\text{P422})'_2$
4_x	$(0, \frac{1}{2}, 0)$	$(0, \frac{1}{2}, \frac{1}{2})$	$(\text{P422})'_3$

Minimal Supergroup Data

$P222$

No. 16

$P222$

I Minimal *translationengleiche* supergroups

[2] $Pmmm$ (47); [2] $Pnnn$ (48); [2] $Pccm$ (49); [2] $Pban$ (50); [2] $P422$ (89) [2] $P4_222$ (93); [2] $P\bar{4}2c$ (112); [2] $P\bar{4}2m$ (111); [3] $P23$ (195)

II Minimal non-isomorphic *klassengleiche* supergroups

- Additional centring translations


[2] $A222$ (21, $C222$); [2] $B222$ (21, $C222$); [2] $C222$ (21); [2] $I222$ (23)

- Decreased unit cell

Incomplete data

Space-group type only

No transformation
matrix



Problem: SUPERGROUPS OF SPACE GROUPS SUPERGROUPS MINSUP

Click here to see the list with all minimal supergroups of a given space group(MINSUP)

supergroup

Please, enter the sequential numbers of group and supergroup as given in the *International Tables for Crystallography, Vol. A*:

Enter supergroup number (G) or choose it:

89

Enter group number (H) or choose it:

16

Enter the index [G:H]

2

space group

index

By default the Euclidean normalizers are used. If you want to use other normalizer, please check it from the list below:

Group normalizer

Euclidean normalizer

Subgroup normalizer

Euclidean normalizer

Subgroup normalizer

Euclidean normalizer

Euclidean normalizer

affine normalizer

user defined normalizer

Output
Supergroups

Find the Supergroups

Supergroups (of index 2) isomorphic to the group 89 (P422)
of the group 16 (P222)

No	Transformation matrix	Coset representatives	Wyckoff Splitting	More...
1	$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$	(x, y, z) (-y, x, z)	[WP splitting]	<input type="button" value="Full cosets"/>
2	$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$	$\begin{bmatrix} 1/2 \\ 0 \\ 0 \end{bmatrix}$ (x, y, z) (-y-1/2, x+1/2, z)	[WP splitting]	<input type="button" value="Full cosets"/>

option
normalizers

ADDITIONAL

ADDITIONAL

GENERATION OF SPACE GROUPS

Generation of space groups

Crystallographic groups are **solvable** groups

Composition series: $P \triangleleft Z_2 \triangleleft Z_3 \triangleleft \dots \triangleleft G$
index 2 or 3

Set of generators of a group is a set of group elements such that each element of the group can be obtained as an ordered product of the generators

$$W = (g_h)^{k_h} * (g_{h-1})^{k_{h-1}} * \dots * (g_2)^{k_2} * g_1$$


g_1 - identity

g_2, g_3, g_4 - primitive translations

g_5, g_6 - centring translations


g_7, g_8, \dots - generate the rest of elements

Generation of sub-cubic point groups




Generation of orthorhombic and tetragonal groups

Hermann–Mauguin symbol of crystal class	Generators G_i (sequence left to right)
1	1
$\bar{1}$	$\bar{1}$
2	2
m	m
$2/m$	$2, \bar{1}$
222	$2_z, 2_y$
$mm2$	$2_z, m_y$
mmm	$2_z, 2_y, \bar{1}$
4	$2_z, 4$
$\bar{4}$	$2_z, \bar{4}$
$4/m$	$2_z, 4, \bar{1}$
422	$2_z, 4, 2_y$
$4mm$	$2_z, 4, m_y$
$\bar{4}2m$	$2_z, \bar{4}, 2_y$
$\bar{4}m2$	$2_z, \bar{4}, m_y$
$4/mmm$	$2_z, 4, 2_y, \bar{1}$



Generation of sub-hexagonal point groups



Generation of trigonal and hexagonal groups

3	3	1
$\bar{3}$	$3, \bar{1}$	
321 (rhombohedral coordinates)	$3, 2_{110}$ $3_{111}, 2_{10\bar{1}}$)	3
312	$3, 2_{1\bar{1}0}$	
$3m1$ (rhombohedral coordinates)	$3, m_{110}$ $3_{111}, m_{10\bar{1}}$)	
$31m$	$3, m_{1\bar{1}0}$	
$\bar{3}m1$ (rhombohedral coordinates)	$3, 2_{110}, \bar{1}$ $3_{111}, 2_{10\bar{1}}, \bar{1}$)	32
$\bar{3}1m$	$3, 2_{1\bar{1}0}, \bar{1}$	
6	$3, 2_z$	6
$\bar{6}$	$3, m_z$	
$6/m$	$3, 2_z, \bar{1}$	
622	$3, 2_z, 2_{110}$	
$6mm$	$3, 2_z, m_{110}$	
$\bar{6}m2$	$3, m_z, m_{110}$	
$\bar{6}2m$	$3, m_z, 2_{110}$	
$6/mmm$	$3, 2_z, 2_{110}, \bar{1}$	622

EXERCISES

Problem 2.30 (A)

Generate the space group C2mm using the selected generators

Compare the results of your calculation with the coordinate triplets listed under General position of the ITA data of C2mm

General Layout: Right-hand page

① CONTINUED

No. 35

Cmm2

② Generators selected (1); $t(1,0,0)$; $t(0,1,0)$; $t(0,0,1)$; $t(\frac{1}{2},\frac{1}{2},0)$; (2); (3)

③ Positions

Multiplicity, Wyckoff letter, Site symmetry	Coordinates				Reflection conditions
	(0,0,0)+	$(\frac{1}{2},\frac{1}{2},0)+$			General:
8 <i>f</i> 1	(1) x,y,z	(2) \bar{x},\bar{y},z	(3) x,\bar{y},z	(4) \bar{x},y,z	$hkl : h+k = 2n$ $0kl : k = 2n$ $h0l : h = 2n$ $hk0 : h+k = 2n$ $h00 : h = 2n$ $0k0 : k = 2n$
4 <i>e</i> <i>m</i> . .	0,y,z	0, \bar{y},z			Special: as above, plus no extra conditions
4 <i>d</i> . <i>m</i> .	$x,0,z$	$\bar{x},0,z$			no extra conditions
4 <i>c</i> . . 2	$\frac{1}{4},\frac{1}{4},z$	$\frac{1}{4},\frac{3}{4},z$			$hkl : h = 2n$
2 <i>b</i> <i>m m</i> 2	0, $\frac{1}{2},z$				no extra conditions
2 <i>a</i> <i>m m</i> 2	0,0,z				no extra conditions

EXERCISES

Problem 2.30 (B)

Generate the space group **P4mm** using the selected generators.

Compare the results of your calculation with the coordinate triplets listed under General position of the ITA data of **P4mm**

Hint: Construct the composition series for the space group **P4mm** in analogy with the composition series of **4mm**

$$\begin{array}{cccccc} & 2_z & & 4_z & & m_x \\ I <\!\!\triangleleft & 2 & <\!\!\triangleleft & 4 & <\!\!\triangleleft & 4mm \\ [2] & & [2] & & [2] & \end{array}$$

Generators selected (1); $t(1,0,0)$; $t(0,1,0)$; $t(0,0,1)$; (2); (3); (5)

Positions

Multiplicity,
Wyckoff letter,
Site symmetry

Coordinates

8	<i>g</i>	1	(1) x,y,z (5) x,\bar{y},z	(2) \bar{x},\bar{y},z (6) \bar{x},y,z	(3) \bar{y},x,z (7) \bar{y},\bar{x},z	(4) y,\bar{x},z (8) y,x,z
4	<i>f</i>	. <i>m</i> .	$x, \frac{1}{2}, z$	$\bar{x}, \frac{1}{2}, z$	$\frac{1}{2}, x, z$	$\frac{1}{2}, \bar{x}, z$
4	<i>e</i>	. <i>m</i> .	$x, 0, z$	$\bar{x}, 0, z$	$0, x, z$	$0, \bar{x}, z$
4	<i>d</i>	. . <i>m</i>	x, x, z	\bar{x}, \bar{x}, z	\bar{x}, x, z	x, \bar{x}, z
2	<i>c</i>	2 <i>m m</i> .	$\frac{1}{2}, 0, z$	$0, \frac{1}{2}, z$		
1	<i>b</i>	4 <i>m m</i>	$\frac{1}{2}, \frac{1}{2}, z$			

EXERCISES

Problem 2.3I (additional)

Generate the space group $P4_2/mbc$ using the selected generators.

Compare the results of your calculation with the coordinate triplets listed under General position of the ITA data of $P4_2/mbc$

Hint: Construct the composition series for the space group $P4_2/mbc$ in analogy with the composition series of $4/mmm$

$$\begin{array}{cccccc} 2_z & & 4_z & & 2_y & \bar{1} \\ I < \textbf{2} < \textbf{4} < \textbf{422} < \textbf{4/mmm} \\ [2] & [2] & [2] & & [2] & \end{array}$$

$P4_2/mbc$

D_{4h}^{13}



$4/mmm$

Tetragonal

No. 135

$P\ 4_2/m\ 2_1/b\ 2/c$

Patterson symmetry $P4/mmm$



Origin at centre ($2/m$) at $4_2/m1n$

Asymmetric unit $0 \leq x \leq \frac{1}{2}; \quad 0 \leq y \leq \frac{1}{2}; \quad 0 \leq z \leq \frac{1}{4}$

Symmetry operations

- | | | | |
|----------------------------|----------------------------|------------------------------------|---|
| (1) 1 | (2) 2 0,0,z | (3) $4^+(0,0,\frac{1}{2})$ | (4) $4^-(0,0,\frac{1}{2})$ |
| (5) $2(0,\frac{1}{2},0)$ | (6) $2(\frac{1}{2},0,0)$ | $0,0,z$ | $0,0,z$ |
| (9) $\bar{1} \ 0,0,0$ | (10) $m \ x,y,0$ | (7) $2(\frac{1}{2},\frac{1}{2},0)$ | (8) $2 \ x,\bar{x}+\frac{1}{2},\frac{1}{4}$ |
| (13) $a \ x,\frac{1}{4},z$ | (14) $b \ \frac{1}{4},y,z$ | $x,x,\frac{1}{4}$ | (11) $\bar{4}^+ \ 0,0,z; \ 0,0,\frac{1}{4}$ |
| | | (15) $c \ x+\frac{1}{2},\bar{x},z$ | (12) $\bar{4}^- \ 0,0,z; \ 0,0,\frac{1}{4}$ |
| | | | (16) $n(\frac{1}{2},\frac{1}{2},\frac{1}{2}) \ x,x,z$ |

Generators selected (1); $t(1,0,0)$; $t(0,1,0)$; $t(0,0,1)$; (2); (3); (5); (9)

Positions

Multiplicity,
Wyckoff letter,
Site symmetry

Coordinates

Reflection conditions

16	<i>i</i>	1	Coordinates				General:
			(1) x, y, z	(2) \bar{x}, \bar{y}, z	(3) $\bar{y}, x, z + \frac{1}{2}$	(4) $y, \bar{x}, z + \frac{1}{2}$	
			(5) $\bar{x} + \frac{1}{2}, y + \frac{1}{2}, \bar{z}$	(6) $x + \frac{1}{2}, \bar{y} + \frac{1}{2}, \bar{z}$	(7) $y + \frac{1}{2}, x + \frac{1}{2}, \bar{z} + \frac{1}{2}$	(8) $\bar{y} + \frac{1}{2}, \bar{x} + \frac{1}{2}, \bar{z} + \frac{1}{2}$	$0kl : k = 2n$
			(9) $\bar{x}, \bar{y}, \bar{z}$	(10) x, y, \bar{z}	(11) $y, \bar{x}, \bar{z} + \frac{1}{2}$	(12) $\bar{y}, x, \bar{z} + \frac{1}{2}$	$hh\bar{l} : l = 2n$
			(13) $x + \frac{1}{2}, \bar{y} + \frac{1}{2}, z$	(14) $\bar{x} + \frac{1}{2}, y + \frac{1}{2}, z$	(15) $\bar{y} + \frac{1}{2}, \bar{x} + \frac{1}{2}, z + \frac{1}{2}$	(16) $y + \frac{1}{2}, x + \frac{1}{2}, z + \frac{1}{2}$	$00l : l = 2n$
							$h00 : h = 2n$
8	<i>h</i>	<i>m</i> ...	$x, y, 0$	$\bar{x}, \bar{y}, 0$	$\bar{y}, x, \frac{1}{2}$	$y, \bar{x}, \frac{1}{2}$	Special: as above, plus no extra conditions
			$\bar{x} + \frac{1}{2}, y + \frac{1}{2}, 0$	$x + \frac{1}{2}, \bar{y} + \frac{1}{2}, 0$	$y + \frac{1}{2}, x + \frac{1}{2}, \frac{1}{2}$	$\bar{y} + \frac{1}{2}, \bar{x} + \frac{1}{2}, \frac{1}{2}$	
8	<i>g</i>	..2	$x, x + \frac{1}{2}, \frac{1}{4}$	$\bar{x}, \bar{x} + \frac{1}{2}, \frac{1}{4}$	$\bar{x} + \frac{1}{2}, x, \frac{3}{4}$	$x + \frac{1}{2}, \bar{x}, \frac{3}{4}$	$hkl : l = 2n$
			$\bar{x}, \bar{x} + \frac{1}{2}, \frac{3}{4}$	$x, x + \frac{1}{2}, \frac{3}{4}$	$x + \frac{1}{2}, \bar{x}, \frac{1}{4}$	$\bar{x} + \frac{1}{2}, x, \frac{1}{4}$	
8	<i>f</i>	2..	$0, \frac{1}{2}, z$	$\frac{1}{2}, 0, z + \frac{1}{2}$	$\frac{1}{2}, 0, \bar{z}$	$0, \frac{1}{2}, \bar{z} + \frac{1}{2}$	$hkl : h+k, l = 2n$
			$0, \frac{1}{2}, \bar{z}$	$\frac{1}{2}, 0, \bar{z} + \frac{1}{2}$	$\frac{1}{2}, 0, z$	$0, \frac{1}{2}, z + \frac{1}{2}$	
8	<i>e</i>	2..	$0, 0, z$	$0, 0, z + \frac{1}{2}$	$\frac{1}{2}, \frac{1}{2}, \bar{z}$	$\frac{1}{2}, \frac{1}{2}, \bar{z} + \frac{1}{2}$	$hkl : h+k, l = 2n$
			$0, 0, \bar{z}$	$0, 0, \bar{z} + \frac{1}{2}$	$\frac{1}{2}, \frac{1}{2}, z$	$\frac{1}{2}, \frac{1}{2}, z + \frac{1}{2}$	
4	<i>d</i>	2.22	$0, \frac{1}{2}, \frac{1}{4}$	$\frac{1}{2}, 0, \frac{3}{4}$	$0, \frac{1}{2}, \frac{3}{4}$	$\frac{1}{2}, 0, \frac{1}{4}$	$hkl : h+k, l = 2n$
4	<i>c</i>	2/ <i>m</i> ..	$0, \frac{1}{2}, 0$	$\frac{1}{2}, 0, \frac{1}{2}$	$\frac{1}{2}, 0, 0$	$0, \frac{1}{2}, \frac{1}{2}$	$hkl : h+k, l = 2n$
4	<i>b</i>	4..	$0, 0, \frac{1}{4}$	$0, 0, \frac{3}{4}$	$\frac{1}{2}, \frac{1}{2}, \frac{3}{4}$	$\frac{1}{2}, \frac{1}{2}, \frac{1}{4}$	$hkl : h+k, l = 2n$
4	<i>a</i>	2/ <i>m</i> ..	$0, 0, 0$	$0, 0, \frac{1}{2}$	$\frac{1}{2}, \frac{1}{2}, 0$	$\frac{1}{2}, \frac{1}{2}, \frac{1}{2}$	$hkl : h+k, l = 2n$